

“Prevalence of Pyogenic Granuloma among Dental Patients in Northern India: An Original Research”

Dr. SharibAbdus Salam¹, Dr. Mobeen Khan², Dr. Ajay Kumar Kushwaha³,
Dr.Hanjala Safi Rayeen⁴, Dr.Priyanka Singh⁵, Dr.Rajendra Kumar⁶

1. Senior lecturer, Department of Periodontics and Implantology, Government Patna Dental College and Hospital, Patna.

2. Senior lecturer, Department of Oral Medicine and Radiology, ChandraDentalCollege and Hospital, SafedabadBarabanki.

3. Assistant professor, Oral and Maxillofacial Surgeon, Prasad Institute of Medical Collage Lucknow.

4. MDS, Department of Oral Pathology and Microbiology.

5. BDS, Junior Resident

6. MD (Pharma), Associate professor, Prasad Institute of Medical Collage Lucknow.

Corresponding author: Dr. SharibAbdus Salam; Senior lecturer, Department of Periodontics and Implantology, Government Patna Dental College and Hospital, Patna.

BACKGROUND: Pyogenic granuloma is a very common, non-neoplastic reactive growth of the oral cavity. Conservative surgical excision is the main line of treatment. Other treatment modalities are cryosurgery, or lasersurgery. Scar formation and recurrence are the main post treatment complications.

MATERIALS AND METHODS: 500 patients were included in this study in which 325 patients were male and 175 patients were female. Patients with 18 to 65 years of age were included in this study. Patients with severe reduced mouth opening were excluded from the study. Collected data was analyzed by using software SPSS 16.

RESULTS: Out of 325 male patients, 36 (11.08%) patients were diagnosed with pyogenic granuloma and out of 175 female patients, 22 (12.57%) patients were diagnosed with pyogenic granuloma.

CONCLUSION: Pyogenic granuloma is more common in female than male.

KEYWORDS: Pyogenic Granuloma; Cryosurgery.

Date of Submission: 28-10-2019

Date of Acceptance: 12-11-2019

I. Introduction

Pyogenic granuloma or granuloma pyogenicum is a common tumor-like growth of the oral cavity or skin considered to be non-neoplastic in nature¹. The accurate etiology and pathogenesis of pyogenic granuloma (PG) is still not obvious although, many researchers mentioned that pyogenic granuloma occurred due to an intensive reaction of localized connective tissue to a little irritating factor². Many etiological factors for pyogenic granuloma have been reported for instance, primary teeth exfoliation, eruption of adult teeth, traumatic factors, hormonal change, gingival inflammation, chronic irritant factor, medications, vascular lesions, defective restoration, food accumulation or strange object and calculus in the gingiva and poor oral hygiene status³. Later, in 1904, Hartzell termed it as granuloma pyogenicum. It is the most common nonneoplastic tumor-like growth of the oral cavity. PG is a reactive tumor which arises in response to various stimuli due to local irritation and traumatic activity of tumor cells and on the rate of cell death. Prevalence of PG is more common in females in the second decade of life, probably due to vascularization effect of estrogen and progesterone⁴. Although PG does not show infiltrative tendency or malignant transformation potential, the recurrence rate after simple excision is comparatively high about 15.8% and requires a re-excision of the lesion in the near future⁵. Diagnosis of pyogenic granuloma (PG) depends on clinical, Histopathological and Radiographical examinations. “peripheral giant cell granuloma”, “hyperplastic gingival inflammation”, “Kaposi’s sarcoma” “peripheral odontogenic fibroma”, “peripheral ossifying fibroma”, “angiosarcoma” and “hemangioma” should be considered in differential diagnosis of pyogenic granuloma⁶. Pyogenic granuloma is a common benign vascular tumour occurring in all ages. Both skin and mucous membranes can be affected. The tumour consists of capillary proliferations, venules and fibromyxoidstroma. The development of a lesion occurs in three stages and bleeding is a common symptom. The tumour can mimic various other vascular lesions, solid tumours, and soft tissue infections. In recent years, targeted tumour therapies have become the most common cause of drug-induced pyogenic granulomas. The backbone of treatment is surgical procedures including laser therapy. New developments in medical drug therapy include topical and systemic beta-adrenergic receptor

antagoniststimolol and propranolol. Drug therapy is an alternative for young children, ocular and periungual pyogenic granuloma⁷.

II. Aim And Objectives

To study the prevalence of Pyogenic Granuloma among Dental Patients in Northern India

III. Materials

- Instruments used:
 1. Plane mouth mirrors and probe.
 2. Tongue depressor.
 3. Cotton swab
 4. Kidney trays.
 5. Cotton holders.
 6. 2x2 inch gauze pieces.
 7. Disposable gloves.
 8. Towel.
 9. Metallic scale.
 10. Big steel tray.
- Additional:
 1. Torch
 2. Clip board.
 3. Indible pencils.

IV. Methodology

500 patients were included in this study in which 325 patients were male and 175 patients were female. Informed consent was obtained from the patients selected for the study. Patients with 18 to 65 years of age were included in this study. Patients with severe reduced mouth opening were excluded from the study. Patients with severe systemic disorders (bleeding disorders, cardiac disorders etc.) were not included in this study. Patients with other oral disorders were not included. Patients with oral malignancies were excluded from the study. Examination of the oral cavity was performed under good illumination using gloves, probe, mirror and other equipment. Diagnosis is made on the basis of clinical examination and history of the disorders.

STATISTICAL ANALYSIS: The data was compiled using Microsoft excel sheet (Windows 2010) and Collected data was analyzed by using software SPSS 16. All results were compared using percentage.



Photograph No:-1. Photograph Showing Armamentarium Used For Clinical Examination

V. Results

TABLE NO. 1. GENDER WISE DISTRIBUTION OF THE PATIENTS

Male	325
Female	175
Total	500

Out of 500 patients, 325 patients were male and 175 patients were female.





Table no. 2. Gender wise distribution of pyogenic granuloma

	male	%	female	%	Total	%
Pyogenic granuloma	36	11.08	22	12.57	58	11.6

Out of 325 male patients, 36 (11.08%) patients were diagnosed with pyogenic granuloma and out of 175 female patients, 22 (12.57%) patients were diagnosed with pyogenic granuloma. Out of 500 patients 58 (11.6%) were diagnosed as pyogenic granuloma. Females shows higher prevalence than males.



VI. Discussion

In this study out of 325 male patients, 36 (11.08%) patients were diagnosed with pyogenic granuloma and out of 175 female patients, 22 (12.57%) patients were diagnosed with pyogenic granuloma. Out of 500 patients 58 (11.6%) were diagnosed as pyogenic granuloma. Females show higher prevalence than males. This study is in line with the study conducted by Alwan et al⁸.

VII. Conclusion

Pyogenic granuloma is more common in female than male. Early diagnosis helps in the betterment in the quality of life with early treatment. Further study is required for better and accurate results with larger sample size. Recurrence rate is very high in the case of pyogenic granuloma.

References

- [1]. Samatha Y, Reddy TH, Jyothirmai, Ravikiran A, Sankar AJ. Management of oral pyogenic granuloma with sodium tetra decyl sulphate. A case series. N Y State Dent J 2013;79:55- 7.
- [2]. Verma PK, Srivastava R, Baranwal H, et al. Pyogenic granuloma hyperplastic lesion of the gingiva: case reports. Open Dent J (2012); Journal homepage: www.mcmed.us/journal/ijacr.
- [3]. Richa Wadhawan, Gaurav Solanki, Arjun Deo Singh, Anubhav Jain pyogenic granuloma of right dorsum of hard palate in a 13 year old girl: a case report. International Journal of Advances in Case Reports, 2(13):824-828 (2015).
- [4]. Jafarzadeh H, Sanatkhan M, Mohtasham N. Oral pyogenic granuloma: A review. J Oral Sci 2006;48:167-75.
- [5]. Kamal R, Dahiya P, Puri A. Oral pyogenic granuloma: Various concepts of etiopathogenesis. J Oral Maxillofac Pathol 2012;16:79-82.
- [6]. Reet Kamal, Parveen Dahiya, Abhiney Puri. Oral pyogenic granuloma: various concepts of etiopathogenesis. Journal of Oral and Maxillofacial Pathology, 16(1): 79-82 (2012).
- [7]. Wollina U, Langner D, Franca K, Gianfaldoni S, Lotti T, Tchernev G. Pyogenic Granuloma – A Common Benign Vascular Tumor with Variable. Open Access Maced J Med Sci. 2017 Jul 25; 5(4):423-426. <https://doi.org/10.3889/oamjms.2017.111>.
- [8]. ALWAN et al., Prevalence and Distribution of Gingival Pyogenic Granuloma in Sulaimani population- Kurdistan Region –Iraq, Biomed. & Pharmacol. J., Vol. 11(1), 105-111 (2018).



FIGURE:- CLINICAL PHOTOGRAPH OF PYOGENIC GRANULOMA

Dr. Sharib Abdus Salam. “Prevalence of Pyogenic Granuloma among Dental Patients in Northern India: An Original Research”. IOSR Journal of Dental and Medical Sciences (IOSR-JDMS), vol. 18, no. 11, 2019, pp 68-71.