

# Role of Vitamin E in Prevention of Oral Cancer:-A Review

MD. ASAD IQUBAL<sup>1</sup>, MOBEEN KHAN<sup>2</sup>, PRABHAKAR KUMAR<sup>3</sup>, ANJANI KUMAR<sup>4</sup>, KRATIKA AJAI<sup>5</sup>

## ABSTRACT

Oral cancer is one of the major global threats to public health. The development of oral cancer is tobacco related mainly. Vitamin-E can inhibit reaction of the tobacco specific nitrosamine which undergoes specific activation, detoxification process. Dietary substitute such as vitamin-E can prevent oral cancer at a very early stage that is in premalignant lesions, in premalignant conditions. Main action of vitamin E includes increase immunity, controls free radicals mediated cell disturbances, maintains membrane integrity, inhibit cancer cell growth, cytotoxicity. Many past studies suggest the role of antioxidant (vitamin-E) in treatment of oral mucosal lesions particularly includes oral leukoplakia, oral lichen planus, oral submucous fibrosis and oral cancer. Vitamin-E as an antioxidant helps in prevention and slow the growth of Head and Neck cancer, improve the effects of cancer chemotherapy and reduce the side effects from both chemotherapy and radiation therapy for cancer patients. As prevention modality use of Vitamin-E may be beneficial for human beings.

**Keywords:** Free radicals, Oral cancer, Premalignant lesions and Premalignant conditions, Vitamin-E

## INTRODUCTION

Discovery of Vitamin-E in 1922 is credited to Herbert Evans and Katherine Scott Bishop when they found that "antisterility factor-x" was essential for reproduction in rats [1]. The pair showed that rats fed a purified diet of casein (18%), cornstarch (544%), lard (15%), butterfat (9%), salts (4%), adequate Vitamin-A (as cod liver oil), Vitamin-B (as yeast), and Vitamin-C (as orange juice) lost their ability to reproduce [2]. Two year later Barnett Sure confirmed Evans and Bishop's observations after performing similar experiments; he called the substance "Vitamin-E" because vitamins A, B, C, and D were then already known to the scientific community [2]. In 1936 Evans first published the chemical formula of Vitamin-E in the Journal of Biological Chemistry [1].

Vitamin-E is the collective term for a family of chemical substances that are structurally related to alpha-tocopherol. Vitamin E occurs naturally in eight different forms: four tocopherols, alpha ( $\alpha$ )-, beta ( $\beta$ )-, gamma ( $\gamma$ ) - and delta ( $\delta$ ) tocopherol and four tocotrienols, alpha-, beta-, gamma- and delta-tocotrienol. All of these forms consist of a chromanol ring with a long aliphatic side chain, bound to the chromanol ring at the second position. Tocotrienols differ from their corresponding tocopherols in that the saturated phytol side chain is replaced with an unsaturated isoprenoid side chain. The Greek characters refer to the number and position of the methyl groups at the 5, 7 and 8 positions [3].

Vitamin-E exhibit antioxidant properties by acting as a lipid-soluble free radical scavenger in cell membranes. Thus, Vitamin-E may involve in both initiation and promotion stages. Among the other potentially anticarcinogenic effects of Vitamin E are its ability to inhibit formation of the carcinogenic chemical nitrosamine from nitrites in some foods, and its ability to promote immune system function [4].

## THE ROLE OF VITAMIN-E IN CANCER CHEMOPREVENTION CAN BE SUMMARIZED AS:

1. Inhibits oral cavity carcinogenesis.
2. Reduces the risk of developing oral cancer.
3. Causes reversal of premalignant lesion like oral leukoplakia.

Oxidative damage is recognized as playing a role in the pathogenesis of cancer which could arise from incorrect nutritional habits and lifestyle practices. This process can cause DNA damage, which is a basic mechanism in cancer induction [5]. Sufficient antioxidative

status is crucial in free radical defence. To reduce the risk of oral and pharyngeal cancer, especially oral cell carcinoma, diet must be optimized, primarily to reduce calorie intake, monosaturated fat and red or processed meat. The important dietary micronutrients that are antioxidant in action include vitamin A,  $\beta$ -carotene, lycopene, Vitamin C, vitamin E (alpha- tocopherol), Zinc and Selenium. Considerable evidence exists suggesting a role for nutrients, particularly the so-called antioxidants vitamin A,  $\beta$ -carotene, vitamin C, vitamin E, lipoic acid, zinc, selenium and spirulina in the prevention of this disease [6]. Antioxidants are group of chemical compounds that can deactivate the free radicals and prevent their formation. Free radicals are oxidants which are single unpaired electron that bombard and destroy cells and other molecules in their search for another electron [7]. A recent study has suggested that these anti-oxidant nutrients act to inhibit the development of cancer cells and to destroy them through apoptosis (programmed cell death), by their stimulation of cytotoxic cytokines, by their action on gene expression, by preventing the development of tumour's necessary blood supply or by cellular differentiation [8]. A report has also shown a reduction in adverse effects of chemotherapy when given concurrently with antioxidants [9].

Tocopherol (AT) is the commonest and most active form of vitamin-E. It is found in plant oil, margarine, and green leaves. Tocopherol is an effective antioxidant at high levels of oxygen, protecting cellular membranes from lipidic peroxidation. Main actions of AT includes;

- Free radical scavenging
- Maintenance of membrane integrity
- Immune function
- Inhibition of cancer cell growth/differentiation
- Cytotoxicity
- In hibits mutagenicity and nitrosamine formation
- In hibition of DNA and RNA, protein synthesis in cancer cells.

## ROLE OF VITAMIN E IN PREVENTION OF ORAL CANCER

Chandra Mouli PE et al., [10] recommended use of Vitamin-E as an antioxidant in oral lesions. Vitamin-E can inhibit reactions of the tobacco specific nitrosamine (carcinogens) which undergo specific activation and detoxification process. Antioxidants such as  $\beta$  carotene, provitamin A, vitamin-C, vitamin-E, zinc, selenium and spirulina are believed to have a preventive role against oral cancer.

Recent studies by Balwant Rai et al., [11] have proved that antioxidants such as Vitamin-C and Vitamin-E may be utilized in oral Lichen planus patients to counteract free radical mediated cell disturbances.

T.N. Uma Maheswari [12] highlighted the role of Vitamin-E in the treatment of leukoplakia. Antioxidant combinations [Vitamin A, E, and C] had proved to be most effective with maximum clinical resolution [90%] recorded [13] and regression of dysplasia recorded as 97.5% [14].

According to Uplabdh Raghuvanshi et al., [15] the low levels of  $\alpha$ -tocopherol in patients could be either a cause or effect of oral carcinoma.

Vikas Fotedar et al., [16] states the role of Vitamin-E in oral cancer and chemoprevention. The term vitamin-E describes a family of light antioxidants, four tocopherols (alpha-, beta-, gamma and delta-). Alpha-tocopherol is the only form of vitamin E that is actively maintained in human body. Vitamin-E (alpha-tocopherol) is a potent antioxidant that neutralizes free oxygen radicals and inhibits carcinogenic nitrosamine formation. They have been reported to block the in vivo formation of N-nitroso compounds and suppress chemical carcinogenesis in animals. They may inhibit cancer development through several mechanisms like stimulation of wild-type p53, down regulation of mutant p53, activation of heat shock proteins, and an antiangiogenic effect mediated by blockage of transforming growth factor-alpha. The antioxidant vitamin E ( $\alpha$ -TF) prevented the development of cancers in oral cavities in animal studies. A phase II study showed that among 43 patients with oral leukoplakia who took 400 IU of vitamin E twice daily for 24wks, 46% had clinical responses and 21% had histological responses. The treatment was well tolerated, without any toxicity, and with good compliance.

Rai Balwant [17] reported in his study, the role of Vitamin-E in oral cancer. Antioxidant defences (vitamin-E and C) are compromised and oxidative stress is increased in patients with oral cancer. A weak antioxidant defense system makes the mucosal cells more vulnerable to the cytotoxic effect of reactive oxygen species. This creates an intracellular environment more favourable for DNA damage and disease progression. So, antioxidant supplement (vitamin C and E) may have role in oral cancer patients.

Sumit Bhateja [18] highlighted the role antioxidants in oral mucosal lesions. The evidence in support of a chemopreventive role for the so called antioxidant nutrients, beta-carotene and Vitamin-E, against oral cavity cancer.

S Chitra and C S Shyamala Devi [19] highlighted the effect of Vitamin-E on oral submucous carcinoma patients. Serum glycoproteins were evaluated in oral squamous cell carcinoma patients treated with radiotherapy and also the effect of vitamin-E on these patients. The levels of glycoconjugates were significantly decreased in radiation treated patients having vitamin-E supplementation. This measurement may be useful in assessing disease progression and identifying patients resistant to therapy and the role of vitamin-E on reduction in glycoconjugate levels of radiation treated oral squamous cell carcinoma patients.

Soma Gupta et al., [20] highlighted the role of Vitamin-E in oral submucous fibrosis. Plasma Vitamin-E level was found to be decreased in grade II and III OSMF cases but not in grade I cases. However, mean Vitamin-E level was found to be decreased ( $9.3 \pm 0.3$  mg/L) as compared to healthy controls (mean =  $10.1 \pm 1.2$  mg/L). Enzymatic antioxidant defense was assessed by SOD activity, which did not show any significant change in any stage of the disease.

Maher et al., [21] evaluated the role of multiple micronutrients consisting of retinol, vitamin-E, vitamin-D, vitamin-B complex and some minerals in the management of OSMF and reported clinical improvement.

Other work has indicated a possibly beneficial role for supplemental vitamin-E in diabetes based on lowered serum levels of glycosylated hemoglobin and triglyceride [22]. Supplemental vitamin-E at a dose of 60-800 ID/day reportedly improves immune function in elderly subjects [23] while a single dose of 1g helped to prevent oxidative damage of DNA [24].

## CONCLUSION

Vitamin-E is an essential nutrient that is receiving growth attention in the prevention of precancerous lesions because of its anti oxidant properties. The main natural sources of vitamin are fresh vegetables, vegetable oils, cereals and nuts. Evidence is increasing that free radical reaction are implicated in the development of degenerative diseases. The body's susceptibility to free radical stress and related damage is associated with the overall balance between the stress level and the antioxidant potential of body tissue. It may be prudent to increase the intake of Vitamin-E and other antioxidant to prevent the body from the increasingly high levels of free radicals derived from the environment and from endogenous sources. Beneficial effect of Vitamin-E in controlling free radical damage in biological system should be most apparent when taken a long term basis.

Oral cancer is generally preceded by precancerous lesions which include leukoplakia, lichen planus, oral submucous fibrosis, oral epithelial dysplasia, erythroplakia. The major risk factor of pharyngeal cancer is tobacco and alcohol in Asian countries chewing tobacco, betel nuts, are major risk factors. In these above lesions Vitamin-E as an antioxidant plays preventive role.

## REFERENCES

- [1] Sen CK, Rink C & Khanna S. Palm oil-derived natural vitamin E alpha-tocotrienol in brain health and disease. *J Am Coll Nutr.* 2010;29:314S-23S.
- [2] Wolf G. The discovery of antioxidant function of vitamin E: the contribution of Henry A. Mattill. *J Nutr.* 2005;135:366.
- [3] Fernando Aguilar, et al. Opinion on mixed tocopherols, tocotrienol tocopherol and tocotrienols as source of vitamin e added as a nutritional substance in food supplements. *The EFSA Journal.* 2008;640:1-34.
- [4] van Poppel G, van den Berg H. Vitamins and Cancer. *Cancer Lett.* 1997;114:195-202.
- [5] Beena Das. Antioxidants in the treatment & prevention of oral cancer. *Kerala Dental Journal.* 2008;31:24-33.
- [6] Devasagayam T, Tilak J, Boloor K, Ghaskadbi S, Lele R. Free radicals and antioxidants in human health: current status and future prospects. *JAPI.* 2004;52:794-804.
- [7] Chapple I. Role of free radicals and antioxidants in the pathogenesis of the inflammatory periodontal diseases. *Clinical Molecular Pathology.* 1996;49:247-55.
- [8] Stephen Hsu, Baldev Singh, George Schuster. Induction of apoptosis in oral cancer cells: agents and mechanisms for potential therapy and prevention. *Oral Oncology.* 2003;39:1-13.
- [9] Sankaranarayanan R, Mathew B, Varghese C, Sudhakaran PR, Menon V, Jayadeep A, et al. Chemoprevention of oral leukoplakia with vitamin A and beta carotene: an assessment. *Oral Oncology.* 1997;33:231-36.
- [10] Chandra Mouli PE, Senthil B, Parthiban S, Malarvizhi AE, Priya R, Subha R. Antioxidants and its role in Oral Cancer- a review. *Indian Journal of Science.* 2012;1:113-15.
- [11] Balwant Rai, Simmi Kharb, Rajnish Jain, SC. Anand. Salivary vitamin E and C in lichen planus. *Gomal J Med Sci.* 2008;6:91-92.
- [12] Uma Maheswari T. N., Treatment of oral leukoplakia with antioxidants- a systematic review. *Int J Pharm Bio Sci.* 2013;4:33-41.
- [13] Kaugars GE, S Silverman Jr. S, Lovas JGL, et al. A clinical trial of antioxidant supplements in the treatment of oral leukoplakia. *Oral Surgery, Oral Medicine, Oral Pathology.* 1994;78: 462-68.
- [14] Barth TJ, Zoller J, Kubler A, S Born IA, and Osswald H, Redifferentiation of oral dysplastic mucosa by the application of the antioxidants beta-carotene, alpha-tocopherol and vitamin C. *International Journal for Vitamin and Nutrition research.* 1997;67:368-76.
- [15] Uplabdh Raghuvanshi, Choudhari SC, Ranjit Patil. Serum alpha-Tocopherol Levels Including Status of Oral Carcinoma Patients. *International Journal of Health Sciences & Research.* 2012;2:60-64.
- [16] Vikas Fotedar, Shailee Fotedar, Rajeev K Seam, Manoj K Gupta. Oral Cancer and Prevention. *International Journal of Pharmaceutical Science Invention.* 2013;2:16-20.
- [17] Rai Balwant. Salivary Levels Vitamin E and C in Different Histological Grading of Oral Cancer. *Pesquisa Brasileira em Odontopediatria e Clinica Integrada.* 2008;8:123-25.
- [18] Sumit Bhateja. Role of antioxidants in oral medicine. *IJPSR.* 2012;3:1971-75.
- [19] Chitra S, Shyamala CS Devi. Effect of vitamin E on protein bound carbohydrate complexes in radiation treated oral squamous cell carcinoma patients. *Indian Journal of Clinical Biochemistry.* 2008;23:92-94.

- [20] Soma Gupta, Reddy MVR and Harinath BC. Role of oxidative stress and antioxidants in aetiopathogenesis and management of oral submucous fibrosis. *Indian Journal of Clinical Biochemistry*. 2004;19:138-41.
- [21] Maher R, Aga P, Johnson NW, Sankaranarayanan R, and Warnakulasuriya S. Evaluation of multiple micronutrient supplementations in the management of oral submucous fibrosis in Karachi, Pakistan. *Nutr. Cancer*. 1997;27:41-47.
- [22] Jain SK, Basu TK, Temple NJ, Garg ML, eds. Oxidative stress, vitamin E and diabetes in Antioxidants in Human Health and Disease. Wallingford, Oxon, UK. *CAB International*. 1999;15:1-64.
- [23] Meydani SN, Meydani M, Blumberg JB, Leka LS, Siber G, Loszewski R, et al. Vitamin E supplementation and in vivo immune response in healthy elderly subjects. *JAMA*. 1997;277: 1380-86.
- [24] Panayiotidis M, Collins AR. Ex vivo assessment of lymphocyte antioxidant status using the comment assay. *Free Radic Res*. 1997;27:533-37.

**PARTICULARS OF CONTRIBUTORS:**

1. Post Graduate Student, Department of Oral Medicine and Radiology, Institute of Dental Sciences, Bareilly, India.
2. Post Graduate Student, Department of Oral Medicine and Radiology, Institute of Dental Sciences, Bareilly, India.
3. Post Graduate Student, Department of Oral Medicine and Radiology, Institute of Dental Sciences, Bareilly, India.
4. Post Graduate Student, Department of Oral Medicine and Radiology, Institute of Dental Sciences, Bareilly, India.
5. Post Graduate Student, Department of Oral Medicine and Radiology, Institute of Dental Sciences, Bareilly, India.

**NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:**

Dr. Mobeen Khan,  
Institute of Dental Sciences, Suresh Sharma Nagar Bareilly, Uttar Pradesh-243006, India.  
Phone : 9455808725, E-mail : drmkhan26@gmail.com

Date of Submission: **Mar 05, 2014**

Date of Peer Review: **Jul 01, 2014**

Date of Acceptance: **Jul 08, 2014**

Date of Publishing: **Oct 20, 2014**

**FINANCIAL OR OTHER COMPETING INTERESTS:** None.